

The ultimate broad spectrum...

MEROMUST

Meropenem Injection 1gm / 500mg

Administration of Meropenem by extended or continuous infusion achieve superior CFR against less-susceptible organisms *Pseudomonas aeruginosa* and *Acinetobacter* species in patients without renal dysfunction.

*Journal of Antimicrobial Chemotherapy (2009)
64, 142–150 doi:10.1093/jac/dkp139*




EFEDRA
Efedra Pharmaceuticals Pvt. Ltd.

Recommended Dose in Adults

Type of Injection	Dose	Dosage Interval
Complicated urinary tract	500 mg	Every 8 hours
Uncomplicated skin and skin structure	500 mg	Every 8 hours
Complicated skin and skin structure	500 mg	Every 8 hours
Gynaecological and Pelvic Inflammatory Disease	500 mg	Every 8 hours
Lower respiratory		
Community-acquired pneumonia	500 mg	Every 8 hours
Nosocomial pneumonia	1 g	Every 8 hours
Complicated intra-abdominal	1 g	Every 8 hours
Meningitis	2 g	Every 8 hours
Septicemia	1 g	Every 8 hours

Dosage in Padiatric Patients

Type of Injection	Dose(mg/kg)	Dosing Interval
Complicated urinary tract	10	Every 8 hours
Uncomplicated skin and skin structure	10 - 20	Every 8 hours
Community-acquired pneumonia	10 - 20	Every 8 hours
Complicated intra-abdominal	20	Every 8 hours
Meningitis	40	Every 8 hours

Meropenem is indicated for

- Lower Respiratory Tract infection
- Complicated urinary tract infections
- Complicated and uncomplicated Skin and Skin Structure
- Complicated intra-abdominal infections
- Gynaecological Infections
- Bacterial Meningitis
- Bacterial Septicemia



Start early